



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ARITHMETIC.

140. Proposed by ALOIS F. KOVARIK, Instructor in Mathematics and Physics, Decorah Institute, Decorah Iowa.

$\frac{1}{7}=0.14285\dot{7}$; $\frac{1}{14}=0.0\dot{7}1428\dot{7}$; $\frac{1}{21}=0.0\dot{0}4761\dot{9}$. Notice that the sum of the figures in each period is equal to 27. This is not true with $\frac{1}{72}$, $\frac{1}{73}$. Is there any general law of which these are special cases, and if so, what is it?

Solution by J. SCHEFFER, A. M., Hagerstown, Md.

When the *period* of a circulating decimal fraction consists of an *even* number of figures, the second half of that period can be found by subtracting each figure of the first half from the figure 9. In this case the sum of the figures constituting a period is always divisible by 9, because we have as many 9's as there are figures in a half-period. Should the half-period have 3 or 6 or 9 or 12, etc., figures, then the sum of the figures in the period will be divisible by 27, and only then. The period of $\frac{1}{72}$ is only 8 and cannot be divisible by 27. The period of $\frac{1}{73}$ consists of eight figures of which the first four are 0, 1, 3, 6, consequently the last four are 9, 8, 6, 3, and the sum of the figures $4 \times 9 = 36$.

141. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pa.

If the alloy in a half-dollar be $\frac{1}{13}$ th of the mass, and the coin be worth a cent if it be all alloy, what should be the exact value of the coin if it be all pure silver?

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.; P. S. BERG, Larimore, N. D.; J. SCHEFFER, A. M., Hagerstown, Md.; and HARVEY M. DAVIS, Brown University, Providence, R. I.

Suppose the half-dollar worth 50 cents.

Then since alloy is worth $\frac{1}{13}$ of a cent, $\frac{1}{13}$ of the mass=silver, is worth $49\frac{1}{13}$ cents.

$\frac{1}{13}$ silver is worth $\frac{1}{13}$ of $49\frac{1}{13}$.

$\frac{1}{13}$ silver is worth $\frac{1}{13}$ of $49\frac{1}{13}$ cents = $54\frac{1}{13}$ cents.

GEOMETRY.

49. Proposed by B. F. FINKEL, A. M., M. Sc., Professor of Mathematics at Springfield, Mo.

Given a conic and two circumscribing triangles of the conic; prove that the six vertices of the triangles are con-conic.

Solution by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, O.

This is most neatly solved, perhaps, by the theory of projection.

Projecting two of the vertices of the first triangle into the circular points